

Epidemiology and Pathogenesis of *Staphylococcus* Bloodstream Infections in Humans: a Review

KLAUDIA LISOWSKA-ŁYSIAK¹ , RYSZARD LAUTERBACH² , JACEK MIĘDZOBRODZKI¹ 
and MAJA KOSECKA-STROJEK^{1*} 

¹Department of Microbiology, Faculty of Biochemistry, Biophysics and Biotechnology,
Jagiellonian University, Kraków, Poland

²Department of Neonatology, Collegium Medicum, Jagiellonian University, Kraków, Poland

Submitted 22 October 2020, revised 18 December 2021, accepted 12 January 2021

Abstract

Staphylococci are among the most frequent human microbiota components associated with the high level of bloodstream infection (BSI) episodes. In predisposed patients, there is a high risk of transformation of BSI episodes to sepsis. Both bacterial and host factors are crucial for the outcomes of BSI and sepsis. The highest rates of BSI episodes were reported in Africa, where these infections were up to twice as high as the European rates. However, there remains a great need to analyze African data for comprehensive quantification of staphylococcal BSI prevalence. The lowest rates of BSI exist in Australia. Asian, European, and North American data showed similar frequency values. World-wide analysis indicated that both *Staphylococcus aureus* and coagulase-negative staphylococci (CoNS) are the most frequent BSI agents. In the second group, the most prevalent species was *Staphylococcus epidermidis*, although CoNS were not identified at the species level in many studies. The lack of a significant worldwide decrease in BSI episodes indicates a great need to implement standardized diagnostic methods and research etiological factors using advanced genetic methods.

Key words: bacteremia, carriage, infection, sepsis, *Staphylococcus*

Introduction

Staphylococcus aureus is one of the most frequently isolated pathogens from the hospital or community-acquired infections. Staphylococci are a large group of bacteria in every environment; however, these bacteria can proliferate only in humans or animals. Many staphylococcal species colonize the skin and mucosal membranes, especially the perineum and pharynx. The other sites that harbor these bacteria are the gastrointestinal tract, vagina, and axilla, but carriage in those areas is less frequent (Kosecka-Strojek et al. 2018). Traditionally, staphylococci have been divided into two groups based on the production of extracellular enzyme coagulase: coagulase-positive staphylococci (CoPS) and coagulase-negative staphylococci (CoNS). The first group is represented by well-known opportunistic pathogens such as *Staphylococcus aureus*, *Staphylococcus schleiferi*, *Staphylococcus intermedius*, and *Staphylococcus pseud-intermedius*, and the second group traditionally includes

nonpathogenic or opportunistic pathogens; however, recently, several clinical reports have presented CoNS as dangerous pathogens, particularly for newborns or immunocompromised patients (Heilmann et al. 2019). A few species, namely *Staphylococcus hyicus*, *Staphylococcus agnetis*, and *Staphylococcus felis*, belong to the third group – coagulase-variable staphylococci. These species are usually grouped with CoPS but cannot produce clumping factors, and coagulase production tests give variable results (Becker et al. 2014). As opportunistic pathogens, staphylococci exhibit saprophytic characteristics under physiological conditions, but the bacteria become severe pathogens under additional infection-facilitating conditions.

Staphylococci are etiological agents of diseases with various localizations, manifestations and/or courses of infection. The most frequent infections are local infections, and the bacteria can cause lesions in various anatomical tissues. Overall, the infections are grouped into skin and soft tissue infections (SSTIs) with

* Corresponding author: M. Kosecka-Strojek, Department of Microbiology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Kraków, Poland, e-mail: maja.kosecka-strojek@uj.edu.pl

© 2021 Klaudia Lisowska-Łysiak et al.

This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

manifestations such as dermatitis, abscesses, furunculosis, boils, folliculitis, impetigo, or mastitis, and also includes other severe diseases such as staphylococcal foodborne disease, toxic shock syndrome, and staphylococcal scalded skin syndrome (SSSS) (Foster 2012; Tong et al. 2015). Staphylococci are also common pathogens of deep tissue infections, including foreign bodies infection. Most studies focused on *S. aureus* infections, but there is strong evidence of the CoNS involvement in severe diseases. Osteomyelitis, otitis, wound infection, endophthalmitis, urinary tract infection, meningitis, or even pneumonia may be caused by *S. epidermidis*, *Staphylococcus saprophyticus*, *Staphylococcus lugdunensis*, and *S. schleiferi* (von Eiff et al. 2002; Becker et al. 2014; Argemi et al. 2019). When staphylococci gain entry into the bloodstream, colonization becomes systemic as bacteremia and then advances to infection.

The literature was screened based on a PubMed search using the terms „staphylococci”, „*Staphylococcus*” and „bloodstream infections” and/or „sepsis”. The publications were then evaluated based on a citation index. Specific criteria were used to describe the worldwide occurrence of *S. aureus*, and CoNS bloodstream infections and/or sepsis, such as: only original articles were included; the data from different geographical regions/countries were analyzed; the articles with the highest number of participants and bacterial strains isolated, and those containing long-term studies or the recent data, were selected to the analysis.

Bacteremia, bloodstream infection, and sepsis

Bacteremia. Bacteremia is characterized by the presence of pathogens in the blood (Pai et al. 2015). Transient bacteremia is limited to one or two days, without any manifestations, and may be caused by some staphylococcal species. Furthermore, the phenomenon does not indicate any further manifestation in healthy hosts (Samet et al. 2006). The presence of bacteria in the blood is eliminated by immunological defense systems and is known in the literature as „natural bacteremia” or „physiological bacteremia”.

Bloodstream infection. However, in predisposed hosts, bacteremia advances to bloodstream infection (BSI), manifesting as an inflammatory response against microorganisms or/and against their metabolites present in the body (Dayan et al. 2016). The BSI can be successfully treated or advances to sepsis (Thomer et al. 2016; Michalik et al. 2020). Sepsis is related to organ dysfunction, perfusion disturbances, or hypotension with accompanying lacticaemia, oliguria, and/or psychological disorders (Samet et al. 2006; Hotchkiss et al. 2016).

Therefore, some *S. aureus* bacteremia complications, such as endocarditis, attributable mortality, embolic

stroke, or recurrent infection during the 12-week follow-up period, are circumstances associated with the increased sepsis frequency from 11% to 43%. When the inflammatory response is triggered by the massive release of pro-inflammatory Th1 cytokines, such as TNF- α , IL-1 β , IL-6, and IFN- γ , a septic shock may occur (Dayan et al. 2016).

Sepsis. Sepsis is the incorrect, inflammatory response of the host organism to infection, and often, it is a result of systemic bloodstream infections. Recently, sepsis is defined as life-threatening organ dysfunction caused by a dysregulated host response to infection (Singer et al. 2016). Untreated sepsis can lead to severe sepsis or septic shock and, consequently, multiple organ failure (Sequential Organ Failure Assessment, SOFA) and death (Stevenson et al. 2016). Sepsis is a critical clinical stage of general toxemia and organ dysfunction, and a patient's inflammatory response interferes with the functioning of vital organs, such as the heart, kidneys, lungs, or liver. Sepsis-3 recommends a new sepsis scoring system, rapid sequential assessment of organ failure (qSOFA), consisting of 3 elements: an altered mental state, respiratory rate, and systolic blood pressure (Minejima et al. 2019). Patients with suspected infection expected to have a prolonged ICU stay can be identified at the bedside with quick SOFA, i.e., alteration in mental status, systolic blood pressure ≤ 100 mmHg, or respiratory rate ≥ 22 /min. Moreover, patients with septic shock can be identified with a clinical construct of sepsis with persisting hypotension requiring vasopressors to maintain MAP ≥ 65 mmHg and having a serum lactate level > 2 mmol/l (18 mg/dl) despite adequate volume resuscitation (Singer et al. 2016).

Pathogenicity of staphylococcal bloodstream infections

Antibiotic resistance. Staphylococci exhibit a wide resistance to antibiotics. One of the most dangerous features of staphylococci is their multi-resistance. Research indicates that both the CoPS and CoNS species have staphylococcal chromosome cassettes *mec* (*SCCmec*) that determine resistance to methicillin in both groups. Furthermore, the *SCCmec* elements of CoNS present extreme diversity, which causes many diagnostic problems (Hosseinkhani et al. 2018). The increase of methicillin-resistant *S. aureus* (MRSA) isolates in hospital and the community due to horizontal gene transfer across bacterial species occurred. The environmental and animal-associated CoNS may be underestimated factor for the spread of the resistance genes into more pathogenic species like *S. aureus* (Kosecka-Strojek et al. 2016; Lisowska-Łysiak et al. 2019). Methicillin and

vancomycin resistance remain the major antimicrobial resistance phenotype of concern. Although still relatively infrequent, multi-resistant CoNS with reduced susceptibility to glycopeptides are emerging pathogens of clinical concern and should be kept in mind in empirical and rational therapy of BSI (Veatch et al. 1990; Natoli et al. 2009). In recent years an emerging spread of linezolid-resistant *Staphylococcus capitis* and *S. epidermidis* strains in Europe was shown (Tevell et al. 2017; Kosecka-Strojek et al. 2020). An increased resistance is the result of antibiotic pressure, which could select resistant clones among staphylococci.

Virulence determinants and other invasion/evasion determinants. Staphylococci exhibit a strong capacity to infect human hosts by using specific strategies to enable the adherence, invasion, persistence, and evasion of the host's immunity mechanisms. However, the infection ability is not similar for all species within the *Staphylococcus* genus. In general, CoNS isolates present lower levels of virulence determinants than CoPS, but the factors involved in colonization support invasion in the host (Becker et al. 2014). It is especially true in extremely immature infants, in whom more than 80% of late-onset sepsis (diagnosed after 72 hours of life) is caused by CoNS (Lauterbach et al. 2016; Wójkowska-Mach et al. 2019). However, despite the relatively low level of virulence, immature infants with sepsis caused by these pathogens present a wide range of clinical symptoms (Lauterbach et al. 2016). It was shown that *S. lugdunensis* was responsible for sepsis and endocarditis on the 1st day of life in a term newborn, which underwent lotus birth (Ittleman and Szabo 2018). In contrast, *S. aureus* strains often exhibit a vast arsenal of toxins and enzymes involved in staphylococcal pathogenesis. Toxins can lead to a weak response of the human organism because they can degrade certain host cells, manipulate the innate and adaptive immune response, and degrade intercellular junctions, contributing to the *S. aureus* proliferation (Oliveira et al. 2018).

Toxins. One of *S. aureus* toxicity mechanisms is damage to host cell membranes caused by hemolysins, bicomponent leukocidins, or phenol-soluble modulins (Herrera et al. 2016). However, it has been proven that CoNS also secrete toxins and enzymes. Based on CoNS strains whole-genome sequencing (WGS) performed by Argemi et al. (2019), the presence of proteases, lipases, and hemolysins genes in *S. epidermidis*, *S. capitis*, and *Staphylococcus caprae* was shown. Moreover, enterotoxin genes in *S. epidermidis* and *Staphylococcus haemolyticus* genomes were shown (Nanoukon et al. 2018; Argemi et al. 2019). Other toxins produced by *S. aureus* are usually infection-specific, such as enterotoxins or toxic shock syndrome toxin. Furthermore, bacteria-host interactions depend on extracellular

enzymes, and the largest group of enzymes includes proteases. This category consists of serine proteases, the metalloprotease aureolysin, and staphopains that are engaged in the evasion of complement-mediated killing, host tissue destruction, immunoglobulin degradation, and deregulation of fibrinolysis (Miedzobrodzki et al. 2002; Sabat et al. 2008; Kalińska et al. 2012; Martínez-García et al. 2018).

Biofilm formation. Biofilm formation is an additional factor associated with CoPS and CoNS infections (Grzebyk et al. 2013; Argemi et al. 2019). Biofilm formation is one of the staphylococcal survival strategies within host organisms. The presence of staphylococcal biofilms is a key factor involved in bacterial resistance to various groups of antibiotics. Bacterial biofilms are defined as communities of bacterial colonies attached to the host surface and surrounded by exopolymeric matrix substances strictly regulated by numerous proteins engaged in the biofilm life cycle. It was shown that biofilms could evade the host immune response, which leads to the persistence of staphylococci. Bhattacharya et al. (2018) proved that *S. aureus* biofilms could release leukocidins, which affect extracellular trap formation and allow evasion of neutrophil-mediated killing. Biofilm production has also been proven for CoNS species, including *S. lugdunensis* that produces adhesins and other biofilm promoters (Argemi et al. 2017). Staphylococcal pathogenesis is a process that involves an array of extracellular proteins, biofilm, and cell wall components that are coordinately expressed in different phases of infection. The expression or suppression of two divergent loci, accessory gene regulator (*agr*) and staphylococcal accessory regulator (*sar*) are recognized as critical regulators of virulence in staphylococci (Arya and Princy 2013).

Risk factors present in humans: predisposed patients. Several studies have shown that host risk factors may significantly enhance the effects of BSIs, including mortality. The high-risk group of staphylococcal infections contains mostly patients with indwelling medical devices. The highly predisposed groups also contain premature newborns or elderly patients or multimorbid, chronically ill, or immunosuppressed patients. A large group of the patients infected is also those with inserted foreign plastic bodies, such as implants and venflons.

The essential patients' factors that determine bloodstream infections and complications are age, presence of comorbidities, and appropriate initial antibiotic treatment (Ayau et al. 2017). Bloodstream infections occur in elderly patients over 75 years old, resulting in increased mortality (Gasch et al. 2013). A 9-year study performed by Ayau et al. (2017) underlined risk factors that increased the probability of mortality, such as age, cancer, heart disease, neurological disease, nursing

home residence, and Charlson score greater than 3. In fact, cancer itself increases the 30-day mortality, but Bello-Chavolla et al. (2018) reported additional risk factors, including hematologic malignancy, hyperglycemia, abdominal source of infection, and endocarditis, based on studies conducted on patients with cancer. Malignancy was also confirmed to be a key factor associated with poor outcomes of infection in other studies (Papadimitriou-Olivgeris et al. 2019).

Epidemiology: Worldwide distribution of staphylococcal bloodstream infections

Staphylococcal bloodstream infections are currently a challenging issue for clinicians, diagnosticians, and microbiologists, primarily due to their high frequency worldwide. Studies on bloodstream infection episodes differ slightly from each other because of the high number of patients and the number of institutions involved in providing the data. Interestingly, all of these studies confirmed a high number of staphylococci isolated from blood samples, ranging from 23.9 to 79.2% (Table I). In many cases, *S. aureus*, usually MRSA isolates, and CoNS were the predominant species involved in BSI episodes. However, most importantly, staphylococcal bloodstream infections affect the whole world, not only developing countries. It is imperative to analyze the data to implement standard diagnostic methods, to compare the results among various countries, to evaluate existing preventive measures, and to plan effective infection prevention and control programs or establish new programs, including the use of advanced genetic methods (Dik et al. 2016; Sabat et al. 2017; Kosecka-Strojek et al. 2019). This study compares staphylococcal bloodstream infections in the world. The evaluation of *S. aureus* and CoNS as etiological agents of BSI of the cited publications was made under the following criteria: the studies included patients with symptoms of BSI/sepsis; pathogens grew on at least one percutaneous blood culture and a culture of the catheter tip; bacteria have been identified as *S. aureus* or CoNS species using commercial/automated identification tests; susceptibility testing was performed, and CoNS species from positive blood samples were included in comprehensive data for analysis except where specified in the laboratory records as contaminants.

Europe. The epidemiology of BSI episodes in Europe was analyzed in detail. The European Centre for Disease Prevention and Control (ECDC 2008; 2018) presented that CoNS were the most numerous bloodstream infections pathogens isolated in Europe. Moreover, the biggest groups of infected patients consisted of neonates and children, and the probability of serious complications such as long-term adverse neurological outcomes

or mortality remained high for these infections (Zingg et al. 2017). Deptuła et al. (2018) reported that catheter-related BSI episodes in Poland occurred in 48.9% of the patients analyzed, and the predominant pathogens were CoNS. These results suggest a strong need for the construction of functional incidence-based surveillance programs in Poland to reduce BSI episodes. The Neonatology Surveillance Network (PNSN) prepared one of these programs and focused on late-onset BSI (LO-BSI) in very-low-birth-weight infants. The study showed that CoNS were the most common cause of LO-BSI (Wójkowska-Mach et al. 2014). Both studies confirmed that it is necessary to implement a national program for infectious disease monitoring and prevention.

Another study in Germany focused on pediatric BSI was based on 20 years of sample collection at a tertiary care hospital. This study conducted a complex observation of a large group of BSI episodes. The results showed an increasing number of CoNS to be responsible for these infections (Hufnagel et al. 2008). Similar results were published by Buetti et al. (2017), which were based on a 7-year surveillance study in Switzerland, although the major pathogen isolated was *Escherichia coli*. These findings were confirmed by other studies performed in Switzerland when staphylococci caused a big group of BSI episodes, but the major isolated pathogens were Gram-negative rods (Papadimitriou-Olivgeris et al. 2019).

On the other hand, an increase in the presence of CoNS was observed, but a minority of studies identified bacteria to the species level. One of these studies was performed in Sweden and showed the CoNS were related to newborns' sepsis from 1987 to 2014. The authors presented that *S. epidermidis* (67.4%) was the most frequent pathogen, followed by *S. haemolyticus* (10.5%), and *S. capitis* (9.6%) (Ehlersson et al. 2017). The epidemiological study in France was partially consistent with previously mentioned research and showed that *E. coli* was the primary pathogen in 36% of BSI episodes, followed by *S. aureus* (16%), and CoNS (8%). The other investigation from France showed the median rate of CoNS in sepsis (12.2%), and all of these strains belonged to *S. capitis* species (Butin et al. 2017). However, studies in the United Kingdom, Greece, Netherlands, and Romania confirmed that CoNS were predominant pathogens in BSI episodes and sepsis (Cailes et al. 2018; Zlatian et al. 2018; Gkentzi et al. 2019; Zonnenberg et al. 2019).

Asia. A study designed in Japan by Takeshita et al. (2017) showed that the major pathogens isolated from BSIs were CoNS (736 cases, 23%), but *S. aureus* isolates were also among the most commonly isolated strains. These results were comparable to those observed in Europe (Takeshita et al. 2017). The authors also focused on 30-day mortality associated with the species and the

Table 1
Worldwide distribution of staphylococcal bloodstream infections. The gray areas consist of the primary pathogen isolated in studies provided according to adequate reference.

No.	Continent	Country	No. of institutions	Years of isolation	Total number of BSI episodes	<i>Staphylo- coccus</i>	<i>Staphylo- coccus</i> (%)	<i>S. aureus</i>	<i>S. aureus</i> (%)	CoNS	CoNS (%)	Reference
1	Africa	Ghana	1	2010–2013	1,763	507	28.8	76	4.3	431	24.4	Labi et al. 2016
2	Africa	Egypt	1	2013–2015	65	26	40.1	6	9.3	20	30.8	Seliem and Sultan 2018
3	Africa	Ethiopia	1	2016–2017	88	38	43.0	16	18.0	22	25.0	Sorsa et al. 2019
4	Africa	Zambia	1	2013–2014	103	13	12.0	6	6.0	7	6.0	Kabwe et al. 2016
5	Asia	Japan	5	2012–2013	3,284	1,030	32.2	294	9.2	736	23.0	Takehita et al. 2017
6	Asia	South Korea	55	2013–2014	717	349	48.7	81	11.3	268	37.4	Lee et al. 2015
7	Asia	Arab States	4	2013–2015	785	289	36.85	17	2.2	272	34.65	Hammond et al. 2017
8	Asia	China	1	2015–2016	133	64	60.3	8	7.5	56	52.8	Jiang et al. 2016
9	Asia	Nepal	1	2017	56	50	89.2	11	19.6	39	69.6	Thapa et al. 2019
10	Asia	India	1	2012–2014	183	87	47.4	42	22.9	45	24.5	Bandyopadhyay et al. 2018
11	Asia	Taiwan	1	2008–2013	2,090	485	23.2	57	2.7	428	20.5	Chen et al. 2016
12	Australia	Australia	23	2008–2012	9,418	3,160	36.4	1,429	18.0	1,731	18.4	Si et al. 2016
13	Australia	Australia	1	2005–2016	203	115	40.3	46	16.1	69	24.2	Worth et al. 2018
14	Australia	Australia	1	2005–2016	146	79	54.1	26	17.8	53	36.3	Gowda et al. 2017
15	Europe	Turkey	1	2003–2009; 2010–2016	925	542	58.6	46	5.0	496	53.6	Mutlu et al. 2019
16	Europe	Switzerland	20	2008–2014	1,823	535	30.0	300	17.0	235	13.0	Buetti et al. 2017
17	Europe	Switzerland	1	2014–2017	404	78	19.3	68	16.8	10	2.5	Papadimitriou-Olivgeris et al. 2019
18	Europe	Poland	nd	2012–2015	329	150	45.6	53	16.1	97	29.5	Deptula et al. 2018
19	Europe	Germany	1	1985–1995; 1997–2006	1,646	650	79.2	241	28.1	409	51.1	Hufnagel et al. 2008
20	Europe	Romania	1	2016–2017	170	81	47.65	63	37.06	18	10.59	Zlatian et al. 2018
21	Europe	Holland	1	2008–2014	93	84	90.4	70	75.3	14	15.1	Zonnenberg et al. 2019
22	Europe	France	1	2011–2012	201	28	12.2	0	0.0	28	12.2	Butin et al. 2017
23	Europe	United Kingdom	30	2005–2014	3,903	2,466	65.0	233	8.0	2,233	57.0	Cailes et al. 2017
24	Europe	Greece	16	2012–2015	459	140	30.4	2	0.4	138	30.0	Gkentzi et al. 2019
25	North America	USA	1	2002–2012	8,196	4,254	51.9	721	8.8	3,533	43.1	Larru et al. 2016
26	North America	USA	1	2006–2017	92	39	42.4	7	7.6	32	34.8	Wagstaff et al. 2019
27	North America	USA	10	2015–2018	5,066	1,500	29.0	1,115	22.0	355	7.0	Khare et al. 2019
28	North America	USA	1	2013–2017	97	29	29.9	17	17.5	12	12.4	Black et al. 2019
29	South America	Brazil	28	2016	47	17	36.2	6	12.8	11	23.4	Braga et al. 2018
30	South America	Latin America	32	2001–2013	3,066	1,625	53.0	267	8.7	1,358	44.3	Escalante et al. 2018

nd – no data

group of pathogens. They concluded that the highest mortality rates were exhibited by hospital-acquired BSI (HA-BSI) pathogens, followed by community-onset healthcare-associated BSI (CHA-BSI), and the most dangerous species were CoNS and *Klebsiella pneumoniae*. The study from South Korea showed that CoNS were the most frequent pathogens engaged in neonatal sepsis (Lee et al. 2015). Studies in India proved a high staphylococci frequency in BSI episodes, however, gram-negative rods were mostly isolated in majority from blood samples (Bandyopadhyay et al. 2018). On the other hand, CoNS became the major isolated pathogen in neonatal sepsis in China, Nepal, Taiwan, Turkey and the Arab States, which proves widespread staphylococcal-caused sepsis in Asia (Jiang et al. 2016; Chen et al. 2017; Hammoud et al. 2017; Thapa and Sapkota 2019; Mutlu et al. 2020). These findings strongly correlate with European data.

North America and South America. In the USA research performed by Larru et al. (2016) presented similar results as European or Asian studies. The most commonly isolated pathogens were CoNS and *S. aureus*, and these pathogens were associated with healthcare-acquired BSI. Moreover, the authors confirmed that all the CoNS were evidenced as pathogens and not as contaminants. Another study from the USA showed a significant majority of *S. aureus* strains involved in neonate sepsis (Khare et al. 2020). The predisposed patients' characteristics were also comparable to those observed in Europe. These studies showed that the most endangered group consists of children and infants, especially with prolonged hospitalization.

Interestingly, children hospitalized since birth exhibited a significantly low prevalence of hospital-onset *S. aureus* bacteremia (Burke et al. 2009). For comparison with the USA, Latin American countries were also analyzed for staphylococcal bloodstream infections. Arias et al. (2017) presented a paper summarizing the results for nine South American countries, from Mexico to Argentina. This study did not evaluate the number of coagulase-negative staphylococci but showed many MRSA strains found in BSI samples from these countries. Notably, the highest number of participants with MRSA-associated BSI was reported in Brazil. Other studies confirmed that the prevalence of intensive-care unit-acquired infections was higher in Brazilian hospitals than in European countries and in the USA (Braga et al. 2018). On the other hand, the highest rates of CoNS (44.3%) were present in NEOCOSUR studies on five Latin American countries: Argentina, Chile, Paraguay, Peru and Uruguay (Escalante et al. 2018).

Africa. The World Health Organization (WHO) reported that, to date, information regarding bloodstream infections in Africa is scarce due to the lack of

research (Bagheri Nejad et al. 2011). However, it was estimated that the incidence of bloodstream infections (up to 14.8%) in developing countries in Africa was up to twice as high as the average European prevalence (7.1%) (ECDC 2008). The studies cited were not correlated with each other with respect to the microbiological data; the major BSI-associated pathogens presented, such as *Pseudomonas aeruginosa*, *E. coli*, *K. pneumoniae*, *Enterobacter* spp., and *S. aureus*, varied among papers (Bagheri Nejad et al. 2011). Besides, the most recent investigations showed a significant increase in CoNS prevalence in BSI episodes in Africa. Labi et al. (2016) showed a high number of positive blood culture samples (21.9%) among neonates, and the significant pathogens were CoNS. Nanoukon et al. (2017) obtained similar results in Benin, where *S. haemolyticus* and *S. epidermidis* were identified as the most frequently isolated pathogens. Similar results were also obtained in Egypt and Malawi (Mashaly and El-Mahdy 2017; Musicha et al. 2017). CoNS were confirmed as a major pathogen isolated in further investigations on smaller groups of patients, mostly neonates in Egypt and Ethiopia (Seliem and Sultan 2018; Sorsa et al. 2019). In contrast, a study from Zambia reports that the most frequent pathogen isolated from neonates with sepsis was *Klebsiella* sp. (Kabwe et al. 2016).

Australia. The rate of healthcare-associated BSIs in Australia is lower than reported elsewhere in the world, which was confirmed by a study in Queensland on 23 public hospitals and by research conducted by the Victorian Healthcare Associated Infection Surveillance System (VICNISS) Coordinating Centre in Victoria (Si et al. 2016; Gowda et al. 2017; Worth et al. 2018). Papers showed that the most frequently reported pathogens responsible for BSI episodes were CoNS, from 18.4 to 24.2%, and *S. aureus*, from 15.2 to 16.1%.

The distribution of all of the aforementioned staphylococcal bloodstream infections is presented in Table I.

***S. aureus* is one of the most frequent bloodstream infection agents**

According to the ECDC report, *S. aureus* is one of the major agents causing bloodstream infections in Europe. Based on the studies conducted in 25 European countries, the ECDC estimated the dynamic changes in *S. aureus* clones associated with BSI episodes. The report states that the *S. aureus* BSI infection mortality rate was 19.4% of the episodes' total number. Moreover, as expected, the MRSA all-cause mortality (24.4%) was higher than that of MSSA infections (17.1%).

***Spa* types related to *S. aureus* from BSI in Europe.** ECDC also estimated 20 of the most frequent MRSA and MSSA *spa* types. The first group included the

5 most frequent *spa* types, namely, t032 (ST22, 17.9%), t003 (ST225, 8.8%), t008 (ST8, 8.4%), t002 (ST5, 7.7%), and t067 (ST125, 4.4%). Interestingly, the significant increase in incidence was related to the multilocus variable number of tandem repeats analysis type (MLVA type) ST22, and this lineage constituted 36% of the top-ranking isolates in 2011. This MRSA clone was first identified in England and was further detected in Ireland, Germany, Hungary, Portugal, and Northern Italy. The fifth most abundant *spa* type t067 was firstly described in Spain (Grundmann et al. 2014). In comparison, the rates of MSSA *spa*-type frequency were lower than those of MRSA isolates, and the 7 most popular types were t091 (ST7, 5.3%), t084 (ST15, 4.7%), t002 (ST5, 4.6%), t015 (ST45, 3.7%), t008 (ST8, 3.7%), t012 (ST30, 3.4%), and t0127 (ST1, 3.2%) (Grundmann et al. 2014). Two *spa* types (t008 and t002) were present in both MRSA and MSSA infections, which was probably a result of the high overall global frequency of these types, according to Ridom SpaServer (www.spaserver.ridom.de).

***Spa* types related to *S. aureus* from BSI in Poland.**

S. aureus, a key pathogen in BSI episodes, was also identified in a study conducted by our group (Ilczyszyn et al. 2016). This study, performed on neonates and children in Poland, showed the most frequent MRSA genotype to be *spa* type t003-CC5, which is consistent with the data presented by ECDC. Among MSSA strains, the most frequent genotypes belonged to the following *spa* types: t091-CC7, t037-CC30, t008-CC8, and t240-CC10. Additionally, some of the observed genotypes exhibited age-related patterns, and the *spa* type t003, *spa*-CC 002, and CC5 were strongly associated with invasive infections in infants and young children (Ilczyszyn et al. 2016).

***Spa* types related to *S. aureus* from BSI outside**

Europe. A study in China, performed for five years, examined *S. aureus* BSI samples and identified the most frequent *S. aureus* *spa* types and virulence factors. According to these data, the most frequent MRSA *spa* type in China was t030/t037, belonging to MLVA type ST239. These isolates also harbored SCCmec III cassette, which represents the hospital-acquired strains, and an *agr* system I. In comparison, the most frequent Chinese MSSA isolates presented the t318 type ST188 and also harbored *agr* I (Liu et al. 2018).

Latin American research divided the most numerous MRSA strains into three clades (A, B, and C) based on phylogenetic reconstruction. Strains in clade A belonged to ST5, ST105, and ST1011, and a majority of these strains harbored the gene cassette SCCmec I or II (HA-MRSA). Clade B consisted of the MLS types ST8, ST88, ST97, and ST72, accompanied by SCCmec IV variants. The last clade included Argentinian strains belonging to ST30 (Arias et al. 2017).

Non-*S. aureus* staphylococci as bloodstream infection agents

Human skin is colonized by various staphylococcal species, although the most invasive is *S. aureus*, followed by *Staphylococcus auricularis*, *S. capitis*, *S. epidermidis*, *S. haemolyticus*, *Staphylococcus hominis*, *S. saprophyticus*, *Staphylococcus simulans* and *Staphylococcus warneri* (Yu et al. 2017). The CoNS are among the most commonly isolated microorganisms from blood samples. Compared to *S. aureus* strains, which are classified as invasive pathogens, the clinical significance of CoNS needs to be proven. It is essential to estimate whether the presence of CoNS represents true bacteremia or sample contamination. Many of the studies conducted have not estimated the real impact of CoNS associated with blood infections, mainly because these species are less frequent overall; have not identified these bacteria at the species level; or have not distinguished the species' differences. However, several studies have shown that CoNS can cause serious bloodstream infections (Grzebyk et al. 2013; Li et al. 2016; Szczuka et al. 2016). Therefore, host-specific capabilities and strain-specific features need to be reconsidered for an improved understanding of the course of every particular infection, as under favorable conditions, CoNS species may become highly pathogenic (Becker et al. 2014).

CoNS as BSI agents in Europe. A study performed in Belgium showed that most isolated bloodstream infection-associated CoNS strains belonged to *S. epidermidis*, and 77% of these strains were identified as methicillin-resistant *S. epidermidis* (MRSE). All of these strains presented resistance to a wide range of antibiotics, especially erythromycin (*ermA*, *ermC*, and *msrA*), aminoglycosides (*aacA-aphD* and *aadC*), tetracycline (*tetK*), and mupirocin (*mupA*). Molecular typing of these strains assigned 85% of the MRSE strains to clonal complex (CC) 2, consisting of the ST2, ST5, ST59, and ST88 MLVA types (Deplano et al. 2016). Another study on *S. warneri* strains from Poland showed their wide range of pathogenicity factors. These strains were able to adhere to host cells, produce biofilms, invade and destroy epithelial cells, which strongly facilitated bacterial persistence (Szczuka et al. 2016). This finding warrants reconsideration of the role of CoNS in bloodstream infections.

CoNS as BSI agents in the USA. In studies conducted in the USA, many CoNS isolates (n = 602) were found in blood samples from years 2013–2014. The most frequently isolated strains belonged to *S. epidermidis*, *S. lugdunensis*, *S. hominis*, and *S. capitis* (Sader et al. 2016). A high number of blood samples was also analyzed in Japan by Yamada et al. (2017), and 314 methicillin-resistant CoNS (MRCoNS) strains were found. Among the Japanese strains, the predominant strains

belonged to *S. epidermidis* (78.6%), *S. haemolyticus* (14.3%), and *S. capitis* subsp. *ureolyticus*. A high number of CoNS-associated BSI episodes and increasing resistance rate should also be confirmation of the danger based on the presence and spread of these bacteria.

Conclusions

Staphylococci are among the most frequent pathogens causing bloodstream infections, which can advance to sepsis and are often observed in patients with indwelling medical devices or neonates. A high number of *S. aureus* and CoNS-related BSI episodes in high-risk patients had evidenced a significant challenge for clinicians. Many institutions widely document BSI episodes, and there has not been a worldwide decrease in these episodes. It is vital to improve existing prevention and control programs based on analysis of the data to implement standard diagnostic methods and conduct research on etiological factors, including via the usage of advanced genetic methods.

ORCID

Klaudia Lisowska-Łysiak <https://orcid.org/0000-0001-8659-9947>

Ryszard Lauterbach <https://orcid.org/0000-0003-1431-6252>

Jacek Międzobrodzki <https://orcid.org/0000-0003-4252-880X>

Maja Kosecka-Strojek <https://orcid.org/0000-0001-8337-6975>

Authors' contributions

JM and MKS brought the idea of the project. KL-Ł and MKS performed the literature research and data analysis. KL-Ł, JM and MKS drafted the work. RL provided clinical consultation of the data. All authors critically revised the work.

Acknowledgments

This project was financed by funds No N19/MNS/000016 (to MKS) of the Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University.

Conflict of interest

The authors do not report any financial or personal connections with other persons or organizations, which might negatively affect the contents of this publication and/or claim authorship rights to this publication.

Literature

Argemi X, Hansmann Y, Prola K, Prévost G. Coagulase-negative staphylococci pathogenomics. *Int J Mol Sci*. 2019 Mar 11;20(5):1215. <https://doi.org/10.3390/ijms20051215>

Argemi X, Hansmann Y, Riegel P, Prévost G. Is *Staphylococcus lugdunensis* significant in clinical samples? *J Clin Microbiol*. 2017 Nov;55(11):3167–3174. <https://doi.org/10.1128/JCM.00846-17>

Arias CA, Reyes J, Carvajal LP, Rincon S, Diaz L, Panesso D, Ibarra G, Rios R, Munita JM, Salles MJ, et al. A prospective cohort multicenter study of molecular epidemiology and phylogenomics of *Staphylococcus aureus* bacteremia in nine Latin American countries. *Antimicrob Agents Chemother*. 2017 Oct;61(10):e00816–17. <https://doi.org/10.1128/AAC.00816-17>

Arya R, Princy SA. An insight into pleiotropic regulators Agr and Sar: molecular probes paving the new way for antivirulent therapy. *Future Microbiol*. 2013 Oct;8(10):1339–1353. <https://doi.org/10.2217/fmb.13.92>

Ayau P, Bardossy AC, Sanchez G, Ortiz R, Moreno D, Hartman P, Rizvi K, Prentiss TC, Perri MB, Mahan M, et al. Risk Factors for 30-day mortality in patients with methicillin-resistant *Staphylococcus aureus* bloodstream infections. *Int J Infect Dis*. 2017 Aug;61:3–6. <https://doi.org/10.1016/j.ijid.2017.05.010>

Bandyopadhyay T, Kumar A, Saili A, Randhawa VS. Distribution, antimicrobial resistance and predictors of mortality in neonatal sepsis. *J Neonatal Perinatal Med*. 2018 Jul 05;11(2):145–153. <https://doi.org/10.3233/NPM-1765>

Becker K, Heilmann C, Peters G. Coagulase-negative staphylococci. *Clin Microbiol Rev*. 2014 Oct;27(4):870–926. <https://doi.org/10.1128/CMR.00109-13>

Bello-Chavolla OY, Bahena-Lopez JP, Garciadiego-Fosass P, Volkow P, Garcia-Horton A, Velazquez-Acosta C, Vilar-Compte D. Bloodstream infection caused by *S. aureus* in patients with cancer: a 10-year longitudinal single-center study. *Support Care Cancer*. 2018 Dec;26(12):4057–4065. <https://doi.org/10.1007/s00520-018-4275-1>

Bhattacharya M, Berends ETM, Chan R, Schwab E, Roy S, Sen CK, Torres VJ, Wozniak DJ. *Staphylococcus aureus* biofilms release leukocidins to elicit extracellular trap formation and evade neutrophil-mediated killing. *Proc Natl Acad Sci USA*. 2018 Jul 10;115(28):7416–7421. <https://doi.org/10.1073/pnas.1721949115>

Black CG, Tavares L, Stachel A, Ratner AJ, Randis TM. Distribution of late-onset neonatal sepsis pathogens differs in inpatient and outpatient settings. *Am J Perinatol*. 2019 Sep;36(11):1136–1141. <https://doi.org/10.1055/s-0038-1675643>

Braga IA, Campos PA, Gontijo-Filho PP, Ribas RM. Multi-hospital point prevalence study of healthcare-associated infections in 28 adult intensive care units in Brazil. *J Hosp Infect*. 2018 Jul;99(3):318–324. <https://doi.org/10.1016/j.jhin.2018.03.003>

Buetti N, Atkinson A, Kottanattu L, Bielicki J, Marschall J, Kronenberg A; Swiss Centre for Antibiotic resistance (ANRESIS). Patterns and trends of pediatric bloodstream infections: a 7-year surveillance study. *Eur J Clin Microbiol Infect Dis*. 2017 Mar;36(3):537–544. <https://doi.org/10.1007/s10096-016-2830-6>

Burke RE, Halpern MS, Baron EJ, Gutierrez K. Pediatric and neonatal *Staphylococcus aureus* bacteremia: epidemiology, risk factors, and outcome. *Infect Control Hosp Epidemiol*. 2009 Jul;30(7):636–644. <https://doi.org/10.1086/597521>

Butin M, Rasigade JP, Subtil F, Martins-Simões P, Pralong C, Freydière AM, Vandenesch F, Tigaud S, Picaud JC, Laurent F. Vancomycin treatment is a risk factor for vancomycin-nonsusceptible *Staphylococcus capitis* sepsis in preterm neonates. *Clin Microbiol Infect*. 2017 Nov;23(11):839–844. <https://doi.org/10.1016/j.cmi.2017.03.022>

Cailles B, Kortsalioudaki C, Buttery J, Pattanayak S, Greenough A, Matthes J, Bedford Russell A, Kennea N, Heath PT; neonIN network. Epidemiology of UK neonatal infections: the neonIN infection surveillance network. *Arch Dis Child Fetal Neonatal Ed*. 2018 Nov;103(6):F547–F553. <https://doi.org/10.1136/archdischild-2017-313203>

Chen CY, Tien FM, Sheng WH, Huang SY, Yao M, Tang JL, Tsay W, Tien HF, Hsueh PR. Clinical and microbiological characteristics of bloodstream infections among patients with hematological malignancies with and without neutropenia at a medical centre in northern Taiwan, 2008–2013. *Int J Antimicrob Agents*. 2017 Mar;49(3):272–281. <https://doi.org/10.1016/j.ijantimicag.2016.11.009>

Dayan GH, Mohamed N, Scully IL, Cooper D, Begier E, Eiden J, Jansen KU, Gurtman A, Anderson AS. *Staphylococcus aureus*: the

- current state of disease, pathophysiology and strategies for prevention. *Expert Rev Vaccines*. 2016 Nov;15(11):1373–1392. <https://doi.org/10.1080/14760584.2016.1179583>
- Deplano A, Vandendriessche S, Nonhoff C, Dodémont M, Roisinienis O. National surveillance of *Staphylococcus epidermidis* recovered from bloodstream infections in Belgian hospitals. *J Antimicrob Chemother*. 2016 Jul;71(7):1815–1819. <https://doi.org/10.1093/jac/dkw086>
- Deptuła A, Trejnowska E, Dubiel G, Wanke-Rytt M, Deptuła M, Hryniewicz W. Healthcare associated bloodstream infections in Polish hospitals: prevalence, epidemiology and microbiology – summary data from the ECDC Point Prevalence Survey of Healthcare Associated Infections 2012–2015. *Eur J Clin Microbiol Infect Dis*. 2018 Mar;37(3):565–570. <https://doi.org/10.1007/s10096-017-3150-1>
- Dik JWH, Poelman R, Friedrich AW, Panday PN, Lo-Ten-Foe JR, Assen S, van Gemert-Pijnen JEW, Niesters HGM, Hendrix R, Sinha B. An integrated stewardship model: antimicrobial, infection prevention and diagnostic (AID). *Future Microbiol*. 2016 Jan;11(1):93–102. <https://doi.org/10.2217/fmb.15.99>
- ECDC. Annual epidemiological report on communicable diseases in Europe 2008. Stockholm (Sweden): European Centre for Disease Prevention and Control; 2008.
- ECDC. Healthcare-associated infections acquired in intensive care units. In: ECDC. Annual epidemiological report for 2016. Stockholm (Sweden): European Centre for Disease Prevention and Control; 2018.
- Ehlersson G, Hellmark B, Svartström O, Stenmark B, Söderquist B. Phenotypic characterisation of coagulase-negative staphylococci isolated from blood cultures in newborn infants, with a special focus on *Staphylococcus capitis*. *Acta Paediatr*. 2017 Oct;106(10):1576–1582. <https://doi.org/10.1111/apa.13950>
- Escalante MJ, Ceriani-Cernadas JM, D'Apremont I, Bancalari A, Webb V, Genes L, Villarreal L, Munoz E, Tapia JL; NEOCOSUR Neonatal Network. Late onset sepsis in very low birth weight infants in the South American NEOCOSUR Network. *Pediatr Infect Dis J*. 2018 Oct;37(10):1022–1027. <https://doi.org/10.1097/INF.0000000000001958>
- Foster AP. Staphylococcal skin disease in livestock. *Vet Dermatol*. 2012 Aug;23(4):e342–e343, e343. <https://doi.org/10.1111/j.1365-3164.2012.01093.x>
- Gasch O, Camoez M, Dominguez MA, Padilla B, Pintado V, Almirante B, Molina J, Lopez-Medrano F, Ruiz E, Martinez JA, et al.; REIPI/GEIH Study Groups. Predictive factors for mortality in patients with methicillin-resistant *Staphylococcus aureus* bloodstream infection: impact on outcome of host, microorganism and therapy. *Clin Microbiol Infect*. 2013 Nov;19(11):1049–1057. <https://doi.org/10.1111/1469-0691.12108>
- Gkentzi D, Kortsalioudaki C, Cailles BC, Zaoutis T, Kopsidas J, Tsolia M, Spyridis N, Siahianidou S, Sarafidis K, Heath PT, et al.; Neonatal Infection Surveillance Network in Greece. Epidemiology of infections and antimicrobial use in Greek Neonatal Units. *Arch Dis Child Fetal Neonatal Ed*. 2019 May;104(3):F293–F297. <https://doi.org/10.1136/archdischild-2018-315024>
- Gowda H, Norton R, White A, Kandasamy Y. Late-onset neonatal sepsis – A 10-year review from North Queensland, Australia. *Pediatr Infect Dis J*. 2017 Sep;36(9):883–888. <https://doi.org/10.1097/INF.0000000000001568>
- Grundmann H, Schouls LM, Aanensen DM, Pluister GN, Tami A, Chlebowicz M, Glasner C, Sabat AJ, Weist K, Heuer O, et al.; ESCMID Study Group on Molecular Epidemiological Markers; European Staphylococcal Reference Laboratory Working Group. The dynamic changes of dominant clones of *Staphylococcus aureus* causing bloodstream infections in the European region: results of a second structured survey. *Euro Surveill*. 2014 Dec 11;19(49):20987. <https://doi.org/10.2807/1560-7917.ES2014.19.49.20987>
- Grzebyk M, Brzychczy-Włoch M, Piotrowska A, Krzyściak P, Heczko PB, Bulanda M. [Phenotypic evaluation of hydrophobicity and the ability to produce biofilm in coagulase-negative staphylococci isolated from infected very-low-birthweight newborns] (in Polish). *Med Dosw Mikrobiol*. 2013;65(3):149–159.
- Hammoud MS, Al-Taiar A, Al-Abdi SY, Bozaid H, Khan A, AlMuhairi LM, Rehman MU. Late-onset neonatal sepsis in Arab states in the Gulf region: two-year prospective study. *Int J Infect Dis*. 2017 Feb;55:125–130. <https://doi.org/10.1016/j.ijid.2017.01.006>
- Heilmann C, Ziebuhr W, Becker K. Are coagulase-negative staphylococci virulent? *Clin Microbiol Infect*. 2019 Sep;25(9):1071–1080. <https://doi.org/10.1016/j.cmi.2018.11.012>
- Herrera A, Vu BG, Stach CS, Merriman JA, Horswill AR, Salgado-Pabón W, Schlievert PM. *Staphylococcus aureus* β -toxin mutants are defective in biofilm ligase and sphingomyelinase activity, and causation of infective endocarditis and sepsis. *Biochemistry*. 2016 May 03;55(17):2510–2517. <https://doi.org/10.1021/acs.biochem.6b00083>
- Hosseinkhani F, Tammes Buirs M, Jabalameli F, Emaneini M, van Leeuwen WB. High diversity in SCCmec elements among multidrug-resistant *Staphylococcus haemolyticus* strains originating from paediatric patients; characterization of a new composite island. *J Med Microbiol*. 2018 Jul 01;67(7):915–921. <https://doi.org/10.1099/jmm.0.000776>
- Hotchkiss RS, Moldawer LL, Opal SM, Reinhart K, Turnbull IR, Vincent JL. Sepsis and septic shock. *Nat Rev Dis Primers*. 2016 Dec 22;2(1):16045. <https://doi.org/10.1038/nrdp.2016.45>
- Hufnagel M, Burger A, Bartelt S, Henneke P, Berner R. Secular trends in pediatric bloodstream infections over a 20-year period at a tertiary care hospital in Germany. *Eur J Pediatr*. 2008 Oct;167(10):1149–1159. <https://doi.org/10.1007/s00431-007-0651-4>
- Ilczyszyn WM, Sabat AJ, Akkerboom V, Szkarlat A, Klepacka J, Sowa-Sierant I, Wasik B, Kosecka-Strojek M, Buda A, Miedzobrodzki J, et al. Clonal structure and characterization of *Staphylococcus aureus* strains from invasive infections in paediatric patients from South Poland: association between age, spa types, clonal complexes, and genetic markers. *PLoS One*. 2016 Mar 18;11(3):e0151937. <https://doi.org/10.1371/journal.pone.0151937>
- Ittleman BR, Szabo JS. *Staphylococcus lugdunensis* sepsis and endocarditis in a newborn following lotus birth. *Cardiol Young*. 2018 Nov; 28(11):1367–1369. <https://doi.org/10.1017/S1047951118001300>
- Jiang Y, Kuang L, Wang H, Li L, Zhou W, Li M. The clinical characteristics of neonatal sepsis infection in Southwest China. *Intern Med*. 2016;55(6):597–603. <https://doi.org/10.2169/internalmedicine.55.3930>
- Kabwe M, Tembo J, Chilukutu L, Chilufya M, Ngulube F, Lukwesa C, Kapasa M, Enne V, Wexner H, Mwananyanda L, et al. Etiology, antibiotic resistance and risk factors for neonatal sepsis in a large referral center in Zambia. *Pediatr Infect Dis J*. 2016 Jul; 35(7):e191–e198. <https://doi.org/10.1097/INF.0000000000001154>
- Kalińska M, Kantyka T, Greenbaum DC, Larsen KS, Władyska B, Jabaiah A, Bogyo M, Daugherty PS, Wysocka M, Jaros M, et al. Substrate specificity of *Staphylococcus aureus* cysteine proteases – Staphopains A, B and C. *Biochimie*. 2012 Feb;94(2):318–327. <https://doi.org/10.1016/j.biochi.2011.07.020>
- Khare R, Kothari T, Castagnaro J, Hemmings B, Tso M, Juretschko S. Active monitoring and feedback to improve blood culture fill volumes and positivity across a large integrated health system. *Clin Infect Dis*. 2020 Jan 2;70(2):262–268. <https://doi.org/10.1093/cid/ciz198>
- Kosecka-Strojek M, Buda A, Miedzobrodzki J. Staphylococcal ecology and epidemiology. In: Savini V, editor. *Pet-to-man travelling staphylococci. A world in progress*. Cambridge (USA): Academic Press; 2018. p. 11–24. <https://doi.org/10.1016/B978-0-12-813547-1.00002-9>

- Kosecka-Strojek M, Ilczyszyn WM, Buda A, Polakowska K, Murzyn K, Panz T, Bialecka A, Kasprowicz A, Jakubczak A, Krol J, et al. Multiple-locus variable-number tandem repeat fingerprinting as a method for rapid and cost-effective typing of animal-associated *Staphylococcus aureus* strains from lineages other than sequence type 398. *J Med Microbiol*. 2016 Dec 16;65(12):1494–1504. <https://doi.org/10.1099/jmm.0.000378>
- Kosecka-Strojek M, Sabat AJ, Akkerboom V, Becker K, van Zanten E, Wisselink G, Miedzobrodzki J, Kooistra-Smid AMD, Friedrich AW. Development and validation of a reference data set for assigning *Staphylococcus* species based on next-generation sequencing of the 16S-23S rRNA region. *Front Cell Infect Microbiol*. 2019 Aug 7;9:278. <https://doi.org/10.3389/fcimb.2019.00278>
- Kosecka-Strojek M, Sadowy E, Gawryszewska I, Klepacka J, Tomasik T, Michalik M, Hryniewicz W, Miedzobrodzki J. Emergence of linezolid-resistant *Staphylococcus epidermidis* in the tertiary children's hospital in Cracow, Poland. *Eur J Clin Microbiol Infect Dis*. 2020 Sep;39(9):1717–1725. <https://doi.org/10.1007/s10096-020-03893-w>
- Labi AK, Obeng-Nkrumah N, Bjerrum S, Enweronu-Laryea C, Newman MJ. Neonatal bloodstream infections in a Ghanaian Tertiary Hospital: are the current antibiotic recommendations adequate? *BMC Infect Dis*. 2016 Dec;16(1):598. <https://doi.org/10.1186/s12879-016-1913-4>
- Larru B, Gong W, Vendetti N, Sullivan KV, Localio R, Zaoutis TE, Gerber JS. Bloodstream infections in hospitalized children: epidemiology and antimicrobial susceptibilities. *Pediatr Infect Dis J*. 2016 May;35(5):507–510. <https://doi.org/10.1097/INF.0000000000001057>
- Lauterbach R, Wilk B, Bocheńska A, Hurkała J, Radziszewska R. Nonactivated protein c in the treatment of neonatal sepsis: a retrospective analysis of outcome. *Pediatr Infect Dis J*. 2016 Sep;35(9):967–971. <https://doi.org/10.1097/INF.0000000000001247>
- Lee SM, Chang M, Kim KS. Blood culture proven early onset sepsis and late onset sepsis in very-low-birth-weight infants in Korea. *J Korean Med Sci*. 2015 Oct;30(Suppl 1):S67–74. <https://doi.org/10.3346/jkms.2015.30.S1.S67>
- Li S, Guo Y, Zhao C, Chen H, Hu B, Chu Y, Zhang Z, Hu Y, Liu Z, Du Y, et al. *In vitro* activities of tedizolid compared with other antibiotics against Gram-positive pathogens associated with hospital-acquired pneumonia, skin and soft tissue infection and bloodstream infection collected from 26 hospitals in China. *J Med Microbiol*. 2016 Oct 18;65(10):1215–1224. <https://doi.org/10.1099/jmm.0.000347>
- Lisowska-Łysiak K, Kosecka-Strojek M, Bialecka J, Kasprowicz A, Garbacz K, Piechowicz L, Kmet V, Savini V, Miedzobrodzki J. New insight into genotypic and phenotypic relatedness of *Staphylococcus aureus* strains from human infections or animal reservoirs. *Pol J Microbiol*. 2019;68(1):93–104. <https://doi.org/10.21307/pjm-2019-011>
- Liu Y, Du F, Liu P, Mei Y, Wan L, Wei D, Xu H, Zhang W. Molecular epidemiology and virulence features of *Staphylococcus aureus* bloodstream isolates in a regional burn center in China, 2012–2016. *Microb Drug Resist*. 2018 Nov;24(9):1354–1360. <https://doi.org/10.1089/mdr.2017.0209>
- Martínez-García S, Rodríguez-Martínez S, Cancino-Díaz ME, Cancino-Díaz JC. Extracellular proteases of *Staphylococcus epidermidis*: roles as virulence factors and their participation in biofilm. *APMIS*. 2018 Mar;126(3):177–185. <https://doi.org/10.1111/apm.12805>
- Mashaly GES, El-Mahdy RH. Vancomycin heteroresistance in coagulase negative *Staphylococcus* blood stream infections from patients of intensive care units in Mansoura University Hospitals, Egypt. *Ann Clin Microbiol Antimicrob*. 2017 Dec;16(1):63. <https://doi.org/10.1186/s12941-017-0238-5>
- Michalik S, Sundaramoorthy N, Murr A, Depke M, Völker U, Bröker BM, Aamot HV, Schmidt F. Early-stage *Staphylococcus aureus* bloodstream infection causes changes in the concentrations of lipoproteins and acute-phase proteins and is associated with low antibody titers against bacterial virulence factors. *mSystems*. 2020 Jan 21;5(1):e00632–19. <https://doi.org/10.1128/mSystems.00632-19>
- Miedzobrodzki J, Kaszycki P, Bialecka A, Kasprowicz A. Proteolytic activity of *Staphylococcus aureus* strains isolated from the colonized skin of patients with acute-phase atopic dermatitis. *Eur J Clin Microbiol Infect Dis*. 2002 Apr;21(4):269–276. <https://doi.org/10.1007/s10096-002-0706-4>
- Minejima E, Delayo V, Lou M, Ny P, Nieberg P, She RC, Wong-Beringer A. Utility of qSOFA score in identifying patients at risk for poor outcome in *Staphylococcus aureus* bacteremia. *BMC Infect Dis*. 2019 Dec;19(1):149. <https://doi.org/10.1186/s12879-019-3770-4>
- Musicha P, Cornick JE, Bar-Zeev N, French N, Masesa C, Denis B, Kennedy N, Mallewa J, Gordon MA, Msefula CL, et al. Trends in antimicrobial resistance in bloodstream infection isolates at a large urban hospital in Malawi (1998–2016): a surveillance study. *Lancet Infect Dis*. 2017 Oct;17(10):1042–1052. [https://doi.org/10.1016/S1473-3099\(17\)30394-8](https://doi.org/10.1016/S1473-3099(17)30394-8)
- Mutlu M, Aslan Y, Aktürk Acar F, Kader Ş, Bayramoğlu G, Yılmaz G. Changing trend of microbiologic profile and antibiotic susceptibility of the microorganisms isolated in the neonatal nosocomial sepsis: a 14 years analysis. *J Matern Fetal Neonatal Med*. 2020 Nov; 33(21):3658–3665. <https://doi.org/10.1080/14767058.2019.1582633>
- Nanoukon C, Affolabi D, Keller D, Tollo R, Riegel P, Baba-Moussa L, Prévost G. Characterization of human type C enterotoxin produced by clinical *S. epidermidis* isolates. *Toxins (Basel)*. 2018 Mar 27;10(4):139. <https://doi.org/10.3390/toxins10040139>
- Nanoukon C, Argemi X, Sogbo F, Orekan J, Keller D, Affolabi D, Schramm F, Riegel P, Baba-Moussa L, Prévost G. Pathogenic features of clinically significant coagulase-negative staphylococci in hospital and community infections in Benin. *Int J Med Microbiol*. 2017 Jan;307(1):75–82. <https://doi.org/10.1016/j.ijmm.2016.11.001>
- Natoli S, Fontana C, Favaro M, Bergamini A, Testore GP, Minelli S, Bossa MC, Casapulla M, Broglio G, Beltrame A, et al. Characterization of coagulase-negative staphylococcal isolates from blood with reduced susceptibility to glycopeptides and therapeutic options. *BMC Infect Dis*. 2009 Dec;9(1):83. <https://doi.org/10.1186/1471-2334-9-83>
- Nejad SB, Allegranzi B, Syed S, Ellis B, Pittet D. Health-care-associated infection in Africa: a systematic review. *Bull World Health Organ*. 2011 Oct 1;89(10):757–765. <https://doi.org/10.2471/BLT.11.088179>
- Oliveira D, Borges A, Simões M. *Staphylococcus aureus* toxins and their molecular activity in infectious diseases. *Toxins (Basel)*. 2018 Jun 19;10(6):252. <https://doi.org/10.3390/toxins10060252>
- Pai S, Enoch DA, Aliyu SH. Bacteremia in children: epidemiology, clinical diagnosis and antibiotic treatment. *Expert Rev Anti Infect Ther*. 2015 Sep 02;13(9):1073–1088. <https://doi.org/10.1586/14787210.2015.1063418>
- Papadimitriou-Olivgeris M, Psychogiou R, Garessus J, Camaret AD, Fourre N, Kanagaratnam S, Jecker V, Nusbaumer C, Monnerat LB, Kocher A, et al. Predictors of mortality of bloodstream infections among internal medicine patients in a Swiss hospital: role of quick Sequential Organ Failure Assessment. *Eur J Intern Med*. 2019 Jul;65:86–92. <https://doi.org/10.1016/j.ejim.2019.05.003>
- Sabat AJ, van Zanten E, Akkerboom V, Wisselink G, van Slochteren K, de Boer RF, Hendrix R, Friedrich AW, Rossen JWA, Kooistra-Smid AMD. Targeted next-generation sequencing of the 16S–23S rRNA region for culture-independent bacterial identification – increased discrimination of closely related species. *Sci Rep*. 2017 Dec;7(1):3434. <https://doi.org/10.1038/s41598-017-03458-6>

- Sabat AJ, Wladyka B, Kosowska-Shick K, Grundmann H, van Dijk J, Kowal J, Appelbaum PC, Dubin A, Hryniewicz W. Polymorphism, genetic exchange and intragenic recombination of the aureolysin gene among *Staphylococcus aureus* strains. BMC Microbiol. 2008;8(1):129. <https://doi.org/10.1186/1471-2180-8-129>
- Sader HS, Farrell DJ, Flamm RK, Streit JM, Mendes RE, Jones RN. Antimicrobial activity of ceftaroline and comparator agents when tested against numerous species of coagulase-negative *Staphylococcus* causing infection in US hospitals. Diagn Microbiol Infect Dis. 2016 May;85(1):80–84. <https://doi.org/10.1016/j.diagmicrobio.2016.01.010>
- Samet A, Bronk M, Sledzińska A, Labon M, Rybak B. [Nosocomial bacteremia] (in Polish). Przegl Epidemiol. 2006;60(1):35–41.
- Seliem WA, Sultan AM. Etiology of early onset neonatal sepsis in neonatal intensive care unit – Mansoura, Egypt. J Neonatal Perinatal Med. 2018 Sep 28;11(3):323–330. <https://doi.org/10.3233/NPM-17128>
- Si D, Runnegar N, Marquess J, Rajmohan M, Playford EG. Characterising health care-associated bloodstream infections in public hospitals in Queensland, 2008–2012. Med J Aust. 2016 Apr;204(7):276. <https://doi.org/10.5694/mja15.00957>
- Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). JAMA. 2016 Feb 23;315(8):801–810. <https://doi.org/10.1001/jama.2016.0287>
- Sorsa A, Früh J, Stötter L, Abdissa S. Blood culture result profile and antimicrobial resistance pattern: a report from neonatal intensive care unit (NICU), Asella teaching and referral hospital, Asella, south East Ethiopia. Antimicrob Resist Infect Control. 2019 Dec;8(1):42. <https://doi.org/10.1186/s13756-019-0486-6>
- Stevenson M, Pandor A, Martyn-St James M, Rafia R, Uttley L, Stevens J, Sanderson J, Wong R, Perkins GD, McMullan R, et al. Sepsis: the LightCycler SeptiFast Test MGRADE®, SepsiT[™] and IRIDICA BAC BSI assay for rapidly identifying bloodstream bacteria and fungi – a systematic review and economic evaluation. Health Technol Assess. 2016 Jun;20(46):1–246. <https://doi.org/10.3310/hta20460>
- Szczuka E, Krzysińska S, Kaznowski A. Clonality, virulence and the occurrence of genes encoding antibiotic resistance among *Staphylococcus warneri* isolates from bloodstream infections. J Med Microbiol. 2016 Aug 01;65(8):828–836. <https://doi.org/10.1099/jmm.0.000287>
- Takeshita N, Kawamura I, Kurai H, Araoka H, Yoneyama A, Fujita T, Ainoda Y, Hase R, Hosokawa N, Shimanuki H, et al. Unique characteristics of community-onset healthcare-associated bloodstream infections: a multi-centre prospective surveillance study of bloodstream infections in Japan. J Hosp Infect. 2017 May;96(1):29–34. <https://doi.org/10.1016/j.jhin.2017.02.022>
- Tevell S, Hellmark B, Nilsson-Augustinsson Å, Söderquist B. *Staphylococcus capitis* isolated from prosthetic joint infections. Eur J Clin Microbiol Infect Dis. 2017 Jan;36(1):115–122. <https://doi.org/10.1007/s10096-016-2777-7>
- Thapa S, Sapkota LB. Changing trend of neonatal septicemia and antibiotic susceptibility pattern of isolates in Nepal. Int J Pediatr. 2019 Feb 06;2019:1–7. <https://doi.org/10.1155/2019/3784529>
- Thomer L, Schneewind O, Missiakas D. Pathogenesis of *Staphylococcus aureus* bloodstream infections. Annual Review of Pathology: Mechanisms of Disease. 2016 May 23;11(1):343–364. <https://doi.org/10.1146/annurev-pathol-012615-044351>
- Tong SYC, Davis JS, Eichenberger E, Holland TL, Fowler VG Jr. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. Clin Microbiol Rev. 2015 Jul;28(3):603–661. <https://doi.org/10.1128/CMR.00134-14>
- Veach LA, Pfaller MA, Barrett M, Koontz FP, Wenzel RP. Vancomycin resistance in *Staphylococcus haemolyticus* causing colonization and bloodstream infection. J Clin Microbiol. 1990;28(9):2064–2068. <https://doi.org/10.1128/JCM.28.9.2064-2068.1990>
- von Eiff C, Peters G, Heilmann C. Pathogenesis of infections due to coagulase-negative staphylococci. Lancet Infect Dis. 2002 Nov;2(11):677–685. [https://doi.org/10.1016/S1473-3099\(02\)00438-3](https://doi.org/10.1016/S1473-3099(02)00438-3)
- Wagstaff JS, Durrant RJ, Newman MG, Eason R, Ward RM, Sherwin CMT, Enioutina EY. Antibiotic treatment of suspected and confirmed neonatal sepsis within 28 days of birth: A retrospective analysis. Front Pharmacol. 2019 Oct 15;10:1191. <https://doi.org/10.3389/fphar.2019.01191>
- Wójkowska-Mach J, Chmielarczyk A, Strus M, Lauterbach R, Heczko P. Neonate bloodstream infections in organization for economic cooperation and development countries: an update on epidemiology and prevention. J Clin Med. 2019 Oct 21;8(10):1750. <https://doi.org/10.3390/jcm8101750>
- Wójkowska-Mach J, Gulczyńska E, Nowiczewski M, Borszewska-Kornacka M, Domańska J, Merritt TA, Helwich E, Kordek A, Pawlik D, Gadzinowski J, et al. Late-onset bloodstream infections of very-low-birth-weight infants: data from the Polish Neonatology Surveillance Network in 2009–2011. BMC Infect Dis. 2014 Dec;14(1):339. <https://doi.org/10.1186/1471-2334-14-339>
- Worth LJ, Daley AJ, Spelman T, Bull AL, Brett JA, Richards MJ. Central and peripheral line-associated bloodstream infections in Australian neonatal and paediatric intensive care units: findings from a comprehensive Victorian surveillance network, 2008–2016. J Hosp Infect. 2018 May;99(1):55–61. <https://doi.org/10.1016/j.jhin.2017.11.021>
- Yamada K, Namikawa H, Fujimoto H, Nakaie K, Takizawa E, Okada Y, Fujita A, Kawaguchi H, Nakamura Y, Abe J, et al. Clinical characteristics of methicillin-resistant coagulase-negative staphylococcal bacteremia in a tertiary hospital. Intern Med. 2017;56(7):781–785. <https://doi.org/10.2169/internalmedicine.56.7715>
- Yu W, Kim HK, Rauch S, Schneewind O, Missiakas D. Pathogenic conversion of coagulase-negative staphylococci. Microbes Infect. 2017 Feb;19(2):101–109. <https://doi.org/10.1016/j.micinf.2016.12.002>
- Zingg W, Hopkins S, Gayet-Ageron A, Holmes A, Sharland M, Suetens C, Almeida M, Asebergiene J, Borg MA, Budimir A, et al.; ECDC PPS study group. Health-care-associated infections in neonates, children, and adolescents: an analysis of paediatric data from the European Centre for Disease Prevention and Control point-prevalence survey. Lancet Infect Dis. 2017 Apr;17(4):381–389. [https://doi.org/10.1016/S1473-3099\(16\)30517-5](https://doi.org/10.1016/S1473-3099(16)30517-5)
- Zlatian O, Balasoiiu A, Balasoiiu M, Cristea O, Docea A, Mitrut R, Spandidos D, Tsatsakis A, Bancescu G, Calina D. Antimicrobial resistance in bacterial pathogens among hospitalised patients with severe invasive infections. Exp Ther Med. 2018 Sep 14;16(6):4499–4510. <https://doi.org/10.3892/etm.2018.6737>
- Zonnenberg IA, van Dijk-Lokkert EM, van den Dungen FAM, Vermeulen RJ, van Weissenbruch MM. Neurodevelopmental outcome at 2 years of age in preterm infants with late-onset sepsis. Eur J Pediatr. 2019 May;178(5):673–680. <https://doi.org/10.1007/s00431-019-03339-2>